



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Simple and Uniform Method of Obtaining Taylor's, Cayley's, and Lagrange's Series.

BY J. C. GLASHAN, *Ottawa, Canada.*

In a short note in the *American Journal of Mathematics*, Vol. I, p. 287, I have given Taylor's Theorem under the form .

$$f(x+a) = \left(1 - \int_0^a da \cdot \frac{d}{dx}\right)^{-1} f x.$$

I find that although this form is excellent as a merely symbolic notation, the operator $\left(1 - \int_0^a d\alpha \cdot \frac{d}{d\dot{x}}\right)^{-1}$ introduces some obscurity in the reasoning if employed in obtaining the theorem. I therefore propose to show the real method adopted in the note and to apply it to obtain Lagrange's Series.

Taylor's Theorem. Let x and a be independent variables, and $fy, f'y, \dots, f^n y$ be continuous from $y = x$ to $y = x + a$.

Since $\frac{d}{da} f(x+a) = \frac{d}{dx} f(x+a) = f'(x+a)$,

$$\begin{aligned}
 \therefore f(x+a) &= fx + \int_0^a da f'(x+a) \\
 &= fx + \int_0^a da \left\{ f'x + \int_0^a da f''(x+a) \right\} \\
 &= fx + \frac{a}{1} f'x + \int_0^a da \int_0^a da f''(x+a) \\
 &= fx + \frac{a}{1} f'x + \frac{a^2}{2} f''x + \left(\int_0^a da \right)^2 f'''(x+a) \\
 &\quad \dots\dots\dots \\
 &\quad \dots\dots\dots \\
 &= fx + \frac{a}{1} f'x + \dots\dots\dots + \frac{a^{n-1}}{(n-1)!} f^{n-1}x + \left(\int_0^a da \right)^n f^n(x+a)
 \end{aligned}$$

Writing x for $x + a_0 + a_1 + \dots$, the theorem becomes

$$fx = f(x - a_0) + \frac{[a]^1}{1} f'(x - a_0 - a_1) + \dots + \frac{[a]^{n-1}}{(n-1)!} f^{n-1}(x - a_0 - a_1 - \dots - a_{n-1}) + R.$$

Let $\{a\}^n \equiv$ the result of retaining only those terms of the expansion of $(a_0 + a_1 + a_2 + \dots + a_{n-1})^n$ of the form

$$Ca_0^\alpha a_1^\beta a_2^\gamma \dots$$

and affecting each term with the sign $(-1)^{n-m}$ where $m = \frac{\alpha}{a} + \frac{\beta}{\beta} + \frac{\gamma}{\gamma} + \dots$, and the theorem may be written

$$fx = f(x - a_0) + \frac{\{a\}^1}{1} f'(x - a_1) + \dots + \frac{\{a\}^{n-1}}{(n-1)!} f^{n-1}(x - a_{n-1}) + R.$$

At the time of the publication of the above theorem in my note entitled *An Extension of Taylor's Theorem*, (Vol. I, p. 287,) I believed it to be new, but the following note published in *Quart. Journ. Math.*, XIV., 53 (two years before mine), shows the theorem had been given long before by Professor Cayley.

"I wish to put on record the following theorem, given by me as a Senate-House Problem, January, 1851.

If $\{\alpha + \beta + \gamma \dots\}^p$ denote the expansion of $(\alpha + \beta + \gamma \dots)^p$, retaining those terms $N\alpha^a\beta^b\gamma^c \dots$ only in which

$$b + c + d \dots \nless p - 1, \quad c + d \dots \nless p - 2, \quad \&c., \quad \&c.,$$

then

$$x^n = (x + \alpha)^n - n\{\alpha\}^1(x + \alpha + \beta)^{n-1} + \frac{1}{2}n(n-1)\{\alpha + \beta\}^2(x + \alpha + \beta + \gamma)^{n-2} - \frac{1}{6}n(n-1)(n-2)\{\alpha + \beta + \gamma\}^3(x + \alpha + \beta + \gamma + \delta)^{n-3} + \&c.$$

The theorem, in a somewhat different and imperfectly stated form, is given, Burg, *Crelle*, t. I. (1826), p. 368, as a generalization of Abel's theorem

$$(x + \alpha)^n = x^n + n\alpha(x + \beta)^{n-1} + \frac{1}{2}n(n-1)\alpha(\alpha - 2\beta)(x + 2\beta)^{n-2} + \frac{1}{6}n(n-1)(n-2)\alpha(\alpha - 3\beta)^2(x + 3\beta)^{n-3} + \&c."$$

Lagrange's Theorem. Let x and a be independent variables, $u = x + a\phi u$, and $f u, f' u, \dots, f^n u$ continuous from $u = x$ to $u = x + a\phi u$.

$$\frac{dfu}{da} = \phi u \frac{dfu}{dx} = \phi u f' u \frac{du}{dx},$$

$$fu = fx + \int_0^a d\alpha \left(\phi v \cdot \frac{dfv}{dx} \right)$$

Integrating by parts,

$$\begin{aligned}\int_0^a d\alpha \left(\phi v \cdot \frac{dfv}{dx} \right) &= a\phi x \cdot f'x - \int_0^a d\alpha \left\{ \alpha \frac{d}{d\alpha} \left(\phi v \cdot \frac{dfv}{dx} \right) \right\} \\ &= a\phi x \cdot f'x + \frac{1}{2} \int_0^a d\alpha \left[\frac{d(\alpha^2)}{d\alpha} \cdot \frac{d}{dx} \left\{ (\phi v)^2 \frac{dfv}{dx} \right\} \right].\end{aligned}$$

Similarly

$$\begin{aligned}\frac{1}{2} \int_0^a d\alpha \left[\frac{d(\alpha^3)}{d\alpha} \cdot \frac{d}{dx} \left\{ (\phi v)^2 \frac{dfv}{dx} \right\} \right] &= \frac{\alpha^2}{2} \frac{d}{dx} \left\{ (\phi x)^2 f'x \right\} \\ &\quad + \frac{1}{2 \cdot 3} \int_0^a d\alpha \left[\frac{d(\alpha^3)}{d\alpha} \left(\frac{d}{dx} \right)^2 \left\{ (\phi v)^3 \frac{dfv}{dx} \right\} \right];\end{aligned}$$

.....
.....

$$\begin{aligned}\frac{1}{n!} \int_0^a d\alpha \left[\frac{d(\alpha^n)}{d\alpha} \left(\frac{d}{dx} \right)^{n-1} \left\{ (\phi v)^n \frac{dfv}{dx} \right\} \right] &= \\ \frac{\alpha^n}{n!} \left(\frac{d}{dx} \right)^{n-1} \left\{ (\phi x)^n f'x \right\} &+ \frac{1}{n!} \int_0^a d\alpha \left[\alpha^n \left(\frac{d}{dx} \right)^n \left\{ (\phi v)^{n+1} \frac{dfv}{dx} \right\} \right].\end{aligned}$$

$$\therefore fu = fx + \frac{a}{1} \phi x \cdot f'x + \frac{a}{2} \frac{d}{dx} \left\{ (\phi x)^2 f'x \right\} + \dots$$

$$+ \frac{\alpha^n}{n!} \left(\frac{d}{dx} \right)^{n-1} \left\{ (\phi x)^n f'x \right\} + \frac{1}{n!} \int_0^a d\alpha \left[\alpha^n \left(\frac{d}{dx} \right)^n \left\{ (\phi v)^{n+1} \frac{dfv}{dx} \right\} \right].$$

This form of the remainder corresponds to that of Taylor's Theorem obtained by integration by parts. (See also xx₁ of the article following this, entitled *Forms of Rolle's Theorem*.) This form can easily be reduced to that obtained by Zolotareff's method, (*Williamson's Integral Calculus*, 3d edition, p. 159); thus:

$$\begin{aligned}\int_0^a d\alpha \left[\alpha^n \left(\frac{d}{dx} \right)^n \left\{ (\phi v)^{n+1} \frac{dfv}{dx} \right\} \right] &= \left(\frac{d}{dx} \right)^n \int_0^a d\alpha \left\{ (\alpha \cdot \phi v)^n \phi v f'v \frac{dv}{d\alpha} \right\} \\ &= \left(\frac{d}{dx} \right)^n \int_a^0 d\alpha \left\{ (\alpha \cdot \phi v + x - v)^n f'v \frac{dv}{d\alpha} \right\} \\ &= \left(\frac{d}{dx} \right)^n \int_x^u dv \left\{ (\alpha \cdot \phi v + x - v)^n f'v \right\}.\end{aligned}$$

P. S.—I find that a form for the remainder in Lagrange's Series was given by Schlömilch, *Liouville*, III₂, 390, (1858.)